



Death and Postmortem Changes

Done by Mohammed El-Ajou

Definition of Death

- Death is a **process**, not a single event — cells die at different times (brain first, then other tissues).
- It is the **irreversible cessation** of all vital functions, mainly:
 - Heart activity (cardiovascular system)
 - Respiration (respiratory system)
 - Higher brain function (neurological system)
- Absence of the 7 life processes:
 - Movement
 - Reproduction
 - Growth
 - Respiration
 - Nutrition
 - Excretion
 - Sensitivity

Types of Death

1. Somatic Death

- Permanent, irreversible death of the whole organism.
- Loss of consciousness, sensory awareness, and voluntary movement.

2. Clinical Death

- Occurs when the heartbeat and breathing stop.
- Lasts about 4 minutes — CPR can potentially revive life during this period.

3. Cellular Death

- Death of tissues and cells, loss of metabolic functions (especially aerobic respiration).
- Different tissues die at different rates:
 - Brain cortex vulnerable after few minutes of oxygen deprivation.

- Connective tissue and muscles can survive hours or days after circulation stops.



4. Brain Death

- Permanent, irreversible loss of brain function due to lack of oxygen (anoxia).
- **Time frame:** Previously thought 3 minutes caused damage; now extended to 7–9 minutes.
- Loss of cerebral cortex function causes **vegetative state**, but brainstem may remain active (spontaneous breathing).

5. Brainstem Death

- Irreversible damage to the brainstem (vital centers & ascending reticular activating system).
- Patient is comatose, unable to breathe spontaneously, requires life support.
- Accepted medically and legally as **somatic death**.
- Patient may not be dead at the cellular level — this allows for organ transplantation.
- Diagnosed by absence of brainstem reflexes in a comatose, ventilator-dependent patient, with no abnormal postures.

Decorticate Posturing



Cleveland Clinic © 2023

Apparent Death

- A **state of suspended animation** that mimics true death.
- Life processes slow down dramatically but do **not terminate**.
- Can occur in conditions such as:
 - Electrocution
 - Hypothermia
 - Sunstroke
 - Drowning
 - Drug overdose (e.g., barbiturates)
 - Head injury
- Differentiated from true death by monitoring with **ECG (heart activity)** and **EEG (brain activity)**.

difference between Cause, Mode, and Manner of Death:

Term	Definition	Examples
Cause of Death	The specific injury or disease that directly leads to death by causing physiological disruption.	Gunshot wound to the head, stab wound to chest, lung cancer, coronary atherosclerosis
Mode/Mechanism of Death	The physiological state or process caused by the cause that leads to death. It explains <i>how</i> death occurs but not the underlying disease.	Coma, cardiac arrest, pulmonary edema, congestive heart failure
Manner of Death	The classification of death based on circumstances and legal context.	Natural, accident, homicide, suicide, undetermined, pending investigation

Example:

- Cause of death: Gunshot wound to the heart
- Mode of death: Massive hemorrhage
- Manner of death: Could be homicide, suicide, accident, or undetermined depending on circumstances

Medico-legal Aspects of Death

The main aims are:

- **Identify the body** if unknown.
- Determine **time of death**.
- Establish **cause of death**.
- Decide if death is **natural or unnatural**.
- If unnatural, determine if **homicidal, suicidal, or accidental**.

Importance of diagnosing death medico-legally:

1. Detect cause of death.
2. Know time of death.
3. Social reasons (legal, family).
4. Organ donation.
5. Recognize apparent death (e.g., suspended animation).

6. Statistical data.
7. Heritage (inheritance) reasons.

Body Changes After Death

- **Initial changes** start biochemically (cell metabolism shifts to autolysis).
- **Visible changes** occur later, important to:
 - Differentiate normal decomposition from signs of unnatural death.
 - Estimate how long death occurred.

Post-mortem signs can be roughly estimated as:

Timing	Signs/Changes
Immediately	Loss of reflexes, permanent stop of respiration & heartbeat
Early	Changes in eyes, pale skin, muscle flaccidity (primary flaccidity), gastric relaxation and regurgitation
Late	Algor mortis (body cooling), livor mortis (blood pooling), rigor mortis (muscle stiffening)
Very Late	Decomposition processes

Details of early eye changes:

- Loss of corneal and light reflexes.
- Pupils mid-dilated, irregular in size/shape (anisocoria).
- Loss of intraocular tension.
- Retinal vessels show blood column fragmentation (“trucking” or “shunting”).



Early Post-Mortem Changes in Eyelids

- Eyelids are usually **incompletely closed**.

- **Tache noire:**
 - Appears where the sclera (white part of the eye) is exposed to air.
 - Characterized by **two black triangular spots** on each side of the cornea.
 - Caused by **drying** of the exposed sclera.



Early Postmortem Changes

1. Muscles

- **Primary flaccidity:** Rapid loss of muscle tone after death; muscles lose tone but may respond to stimulation for hours.
- **Focal twitching:** Discharges from dying motor neurons can cause small muscle twitches, which decrease over time.
- **Sphincter relaxation:** Loss of muscle tone in sphincters causes urine voiding, useful in estimating time since death.

2. Skin

- Pale skin, conjunctivae, and mucous membranes due to blood pressure drop and circulation cessation.
- Hair follicles die simultaneously with skin; no actual hair growth after death, though beard may appear more prominent.
- Face and lips may remain red or blue in hypoxic or congestive deaths.

3. Stomach

- Gastric contents often found in mouth or airways (~25% of autopsies).
- Regurgitation common during terminal collapse or resuscitation attempts.
- Presence of stomach contents in airways alone does not prove inhalation as cause of death without further evidence.

Algor Mortis (Body Cooling)

- After death, body stops generating heat; temperature gradually approaches ambient temperature.
- Normal oral temperature: 36–37.2°C; rectal: 36.5–37.5°C.
- Rectal temperature <24°C suggests death; <21°C is definitive.
- Cooling rate approx. 1°C/hr in summer, 1.5°C/hr in winter.
- Simple formulas for time since death:
 - Time since death (hours) = $37^{\circ}\text{C} - \text{rectal temperature } (^{\circ}\text{C}) + 3$
 - Time since death = $(98.6^{\circ}\text{F} - \text{rectal temperature } (^{\circ}\text{F})) \div 1.5$
- Assumptions: normal temperature at death and linear cooling may not always hold.
- Factors affecting cooling: body surface area, weight, clothing, posture, environment, infections.

Livor Mortis (Hypostasis)

- Blood circulation stops; gravity causes blood to settle in dependent body parts.
- Red blood cells sediment, causing bluish-red discoloration (“hypostasis”) visible on skin.
- Plasma also settles, contributing to dependent edema and skin blistering.
- Distribution depends on body position:
 - **Supine:** Chest, upper limbs affected.



- **Vertical/Hanging:** Blood pools in lower extremities.



- **Face-down:** Whitening around nose and lips, typical in seizures or intoxication deaths.
- **Common external sites:**
 - Most commonly found on the **back of the shoulders and neck**.
 - May also appear on the **front of the chest**, even if the body is lying on its back.
 - More frequent in **cyanotic** (bluish discoloration due to low oxygen) and **congestive** deaths (e.g., heart failure).
- **Internal lividity (internal discoloration) sites that may mimic disease:**
 - **Heart:** May be mistaken for myocardial infarction (heart attack).
 - **Lungs:** Can resemble pneumonia.
 - **Intestine:** May look like hemorrhagic infarction (localized tissue death due to bleeding).



Color of Hypostasis (Livor Mortis)

- **Variable color** depending on the oxygenation status of the blood at the time of death:
 - **Bright red or pink** in cases of **carbon monoxide poisoning** or **cyanide poisoning** (due to high oxygenation).
 - **Purple or dark red/blue** in cases of **poor oxygenation** (e.g., congestive heart failure, asphyxia).
- May be **masked or obscured** by:
 - Dark skin pigmentation
 - Jaundice (yellowish skin coloration)
 - Certain dermatological conditions affecting the skin

Here's a clear table summarizing the **color of hypostasis** and their causes:

Colour	Cause
Dark blue-pink	Cyanide poisoning
Brown	Methemoglobinemia
Cherry pink	Carbon monoxide (CO) poisoning or cold temperatures
Pale	Anemia, hemorrhage, or normal in extremes of age

Timing and Permanence of Hypostasis

- **Onset:** Begins between 30 minutes to several hours after death.

- **Duration:**
 - In **summer**: lasts 24–36 hours.
 - In **winter**: lasts about 48 hours or more.
 - In **snow**: can last much longer.
- **Effect of Movement:** If the body is moved after death, hypostasis may:
 - Remain fixed in the original position,
 - Move completely to a new position,
 - Or partially move and partially stay fixed.

Medico-Legal Importance of Hypostasis

1. **Sure sign of death** — confirms that death has occurred.
2. **Provides clues about the cause of death.**
3. **Indicates the body's position before and after death.**
4. **Shows if the body was moved after death.**

Here's a clear comparison table for **Hypostasis vs Bruises** based on your information:

Feature	Hypostasis	Bruises
Location	Dependent areas (lowest parts)	Anywhere on the body
Edges	Well-defined	Ill-defined
Blood in Capillaries	Blood retained in intact capillaries	Blood escapes through ruptured capillaries
Surface Level	Same level on the surface	Raised
Color Over Pressure	Pale over pressure areas	Red
Incision Response	Blood flows from the cut vessel (washable)	Blood coagulates in tissue



Mechanism of Rigor Mortis

- **Definition:**

Rigor mortis is a temperature-dependent, physicochemical change in muscle cells after death, causing the muscles to stiffen and limbs to become difficult to move.

- **Process:**

- Death leads to cessation of respiration.
- Oxygen depletion occurs, reducing ATP production.
- Secondary anoxic processes begin, causing lactic acid buildup.
- Acidic cytoplasm and low ATP levels cause actin and myosin filaments in muscle cells to bind tightly, forming a gel-like state.
- This binding prevents muscle relaxation, leading to permanent muscular contraction (rigidity).
- Over time, digestive enzymes break down muscle tissue during decomposition, leading to eventual resolution of rigor mortis.

- **Timing:**

- Starts when ATP levels fall to about 85% of normal.
- Begins approximately **2-3 hours after death**.
- First noticed in smaller muscles (around eyes, mouth, jaw, fingers).
- Peaks within **6-12 hours** after death.
- Resolves around **36-48 hours** postmortem, in the order it developed (called "secondary flaccidity").



Factors Affecting Timing of Rigor Mortis

1. Environmental Temperature

- a. Cold and wet conditions → slower onset, longer duration (rigor lasts **36–48 hours**)
- b. Hot and dry conditions → faster onset, shorter duration (rigor lasts **24–36 hours**)

2. Muscular Activity Before Death

- a. Muscles healthy, robust, and at rest before death → slower onset
- b. Muscles exhausted or fatigued (e.g., from struggling, exertion) → rapid onset, especially in the active limbs
- c. Increased muscular activity before death (e.g., convulsions, electrocution, lightning) → very rapid onset and shorter duration

3. Age

- a. Extremes of age (infants and elderly) → rapid onset

4. Health Status

- a. Poor health or emaciation → delayed or weak rigor mortis
- b. Infants → very rapid onset

5. Other Factors

- a. Certain poisons (e.g., strychnine) causing convulsions can accelerate rigor mortis development

Here's a clear table summarizing **Body Temperature, Muscle Tone, and Estimated Time of Death (ETD)**:

Body Temperature	Muscle Tone	Estimated Time of Death (ETD)
Warm	Flaccid	Less than 3 hours
Warm	Stiff	3 to 8 hours
Cold	Stiff	8 to 36 hours
Cold	Flaccid	More than 36 hours

Rigor Mortis (RM) Specifics:

- **RM in iris:** Can cause unequal rigor, leading to unequal pupil sizes.

- **RM in heart:** Contracted and stiff left ventricle (LV) may be mistaken for LV hypertrophy.
- **RM in dartos muscle (scrotum):** Causes constriction of testes and epididymis; may cause semen expulsion.
- **RM in erector pili muscles (attached to hair follicles):** Causes goosebumps; hair stands up.

Testing for Rigor Mortis:

- Use **gentle pressure** with one or two fingers across a joint.
- Aim: Detect stiffness extent, **do not break the rigor**.
- If rigor is broken by too much force, that muscle group can no longer be reliably tested.

Conditions Mistaken for Rigor Mortis:

1. **Heat Stiffness:**
Exposure to intense heat (burning, electrocution) causes coagulation of muscle proteins → muscle shortening.
2. **Cold Stiffness:**
Exposure to extreme cold (< -5°C) causes solidification of fat and muscles, freezing synovial fluid in joints.

Cadaveric Spasm (Instantaneous Rigor):

- A rare form of muscle stiffening occurring **at the moment of death**, persisting into rigor mortis.
- Usually linked to **violent deaths with intense emotion**.
- May affect whole body muscles or just specific groups (e.g., forearms or hands).
- Common in drowning victims holding onto objects (grass, weeds, roots), showing proof of life at time of water entry.
- Important forensic clue indicating the last action before death.

Here's a clear comparison table between **Rigor Mortis** and **Cadaveric Spasm** based on your details:

Feature	Rigor Mortis	Cadaveric Spasm
Onset	Delayed after death (2-3 hours)	Instantaneous at the moment of death
Duration	Lasts up to 36 hours	Lasts a few hours, then replaced by rigor mortis
Intensity	Moderate intensity	Very strong intensity
Mechanism of formation	Breakdown of ATP below critical level	Unknown; associated with emotions and muscle contraction at death
Muscle involvement	Gradual involvement of all muscles	Selected muscles in contraction at time of death

Decomposition

Decomposition is a mixed process involving:

- **Autolysis:** Internal chemical breakdown of cells by enzymes after death.
- **External decomposition:** Caused by bacteria and fungi from intestines and the environment.
- **Animal scavengers:** Including maggots and mammals contribute to destruction.
- Decomposition varies by body, environment, and even different parts of the same corpse.

Types of Decomposition:

1. Putrefaction

- The normal final sign of death, starting immediately after death on a cellular level.
- Becomes visible within 48–72 hours.
- Two main processes:
 - **Autolysis:** Breakdown by digestive enzymes; accelerated by heat, slowed by cold or freezing. Organs rich in enzymes (e.g., pancreas) autolyze faster than others (e.g., heart).

- **Bacterial action:** Bacteria, mostly from bowel (*Clostridium* species), ferment soft tissues.
- First visible sign: Green or greenish-red discoloration of the anterior abdominal skin, starting usually at the right iliac fossa (24-36 hours).
- Generalized bloating occurs within 60–72 hours.
- Early signs: Yellow-clear fluid-filled blisters, progressing to green-brown fluid-filled blisters.
- Blood vessels act as channels for bacteria, causing "marbling" (linear brown discoloration).
- Skin blisters form and burst, causing sloughing.
- Gas formation causes swelling (face, abdomen, breasts, genitals).
- Eyes and tongue may protrude; bloody fluid (purge) leaks from mouth and nose.
- After about a week, body cavities burst; tissues liquefy and drain.
- Brain and epithelial tissues decompose first; heart, uterus, and prostate decompose more slowly.
- Organ weight decreases as decomposition proceeds.

2. Adipocere

- Chemical change of body fat into a waxy, soap-like substance.
- Requires moisture and bacterial activity (*Clostridium perfringens*).
- Common in wet, warm environments (or from original body water).
- Occurs mainly in subcutaneous fat of cheeks, breasts, buttocks, and sometimes internal organs (liver, kidney, heart).
- Takes months to develop and often partial.
- Stages:
 - Pale, rancid, greasy semi-fluid with unpleasant smell.
 - Becomes brittle and whiter.
 - Fully formed: gray, firm, waxy compound maintaining body shape.
- **Medico-legal importance:**
 - Preserves body allowing identification.
 - May suggest cause of death.
 - Indicates time since death of weeks to months.

3. Skeletonization

- Environment influences this more than time.
- Timeline:
 - 12–18 months: soft tissues absent.
 - 3 years: bones bare and disarticulated.
 - 40–50 years: bones dry, brittle; marrow cavity empty.
- In temperate zones, bones remain solid with marrow preserved for DNA analysis.

4. Mummification

- Occurs in dry, warm conditions (climatic or microenvironment).
- Body dries and shrivels, turning brown.
- No microorganism growth.
- Partial mummification.
- **Medico-legal importance:**
 - Can help determine cause of death.
 - Can detect abnormal pathology in deep tissues.

Determination of Time of Death

1. Core Body Temperature

- a. Most reliable and commonly used method.
- b. Body temperature gradually cools after death until it matches the environment.

2. Rigor Mortis

- a. The stiffening of muscles after death; timing of onset and resolution helps estimate time since death.

3. Hypostasis (Livor Mortis)

- a. Blood settling occurs after death; complete after about 6 hours.

4. Chemical Changes in the Vitreous Humor

- a. Changes in the fluid of the eye can be measured to estimate time since death.

5. Eye Pressure

- a. Eyeballs become softer with less fluid pressure within the first 3 hours after death.

6. Gastric Emptying

- a. The stomach empties depending on meal type and emotional status before death; timing helps estimate death time.

7. Entomology of the Dead

- a. Study of insects and maggots colonizing the body. Different insects appear at various postmortem stages.

8. Scene Markers (Unscientific but often practical)

- a. Uncollected mail or newspapers.
- b. Whether lights are on or off.
- c. How the person is dressed.
- d. Presence of food or dirty dishes.
- e. Sales receipts or dated papers in pockets.
- f. Observations from neighbors about last seen time or changes in habits.